



**PATIENT
EDUCATION**

Bowel Preparation Instructions: MoviPrep

It is very important to follow all the instructions to prepare for your procedure. Your colon must be empty, so the provider can see during the procedure. Many patients feel that the bowel preparation is the hardest part of the procedure.

Preparing for the procedure

- The day before your procedure, you will need to start preparing your colon. Do not eat any solid foods the entire day before your procedure.
- You will be using the low-volume polyethylene glycol (PEG) MoviPrep to clean out your colon.
- Please follow all instructions carefully. If you do not follow the directions properly, then your procedure may be canceled.

Items to buy

- To get ready for your procedure, you will need to buy items for the bowel preparation.
- Be sure to check with your insurance company to see if this bowel preparation is covered.

Prescription required:

- Check with your pharmacy and fill your prescription for MoviPrep preparation kit.

Medication information

Some of the medicines you take may need to be stopped or adjusted temporarily before your procedure.

- **Blood thinners**
 - such as Coumadin (warfarin), Ticlid (ticlopidine hydrochloride), Agrylin (anagrelide), Xarelto (Rivaroxaban), Pradaxa (Dabigatran), Pletal (Cilostazol), Brillinta (Ticagrelor), Eliquis (Apixaban), Effient (Prasugrel), and Plavix (Clopidogrel).
 - You must speak with your prescribing provider or the specialist at least 2 weeks before the scheduled procedure. Do not stop these medications without the consent of your provider.
- **Insulin and diabetes medications**
 - Please call the provider that monitors your glucose levels. Your medications may need to be adjusted due to the diet restrictions required with this bowel preparation. Do not stop these medications without the consent of your provider.
- **Aspirin**
 - You may take your Aspirin as prescribed with a sip of water.

Five (5) days before your procedure

- Do not take medications that will stop diarrhea.
 - Examples: Imodium or Pepto-Bismol.
- Do not take fiber medications or supplements.
 - Examples: Metamucil or Citrucel.

- Do not take any vitamins or supplements.
 - Example: Multi-vitamins or fish oil.

Three (3) days before your procedure

- Begin a low fiber diet. A low fiber diet helps make the cleanout more effective.
- Low fiber examples include:
 - White bread, white rice, pasta, crackers, fish, chicken, eggs, ground beef, creamy peanut butter, cooked/boiled vegetables, canned fruit, bananas, melons, milk, plain yogurt, cheese, salad dressing, and other condiments.
- Do not eat high-fiber foods such as:
 - Popcorn, seeds (flax seeds, sunflower, and quinoa), multigrain breads or pasta, nuts, granola, salads, raw vegetables, or fresh and dried fruits. Limit the amount of meats.

Two (2) days before your procedure

- Increase your water intake.
- Continue a low fiber diet.

Morning check-in

(6:30 a.m. to 11:45 a.m.)

One (1) day before your procedure

- Start your clear liquid diet.
- Only drink clear liquids the entire day before your procedure.
- A clear liquid diet includes:
 - Water, black coffee or tea (no milk or cream), clear broth or bouillon, Gatorade, carbonated or non-carbonated soft drinks (Coke, Sprite), strained fruit juices without pulp (apple, white grape, white cranberry), Jell-O, and popsicles. **Remember, NO RED OR PURPLE LIQUIDS.**
- Do not eat any solid foods.
- If preferred, mix MoviPrep ahead of time and refrigerate before drinking. Please note that MoviPrep mixture should be used within 24 hours.
- At 5:30 p.m., begin to prepare the first dose of the MoviPrep mixture. Empty 1 pouch A and 1 pouch B into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve.
- At 6:00 p.m., start drinking the MoviPrep mixture. Drink an eight-ounce glass of the prep mixture every 15 minutes. You will drink a total of one liter or the entire container of the MoviPrep mixture.
- You will need to finish drinking your prep mixture by 7:00 p.m.
- You will need to be near a bathroom once you start drinking your prep mixture. Some people may feel nauseous, bloated, or vomit. If this happens, take a 15-30-minute break, and then try drinking the prep mixture.
- After you finish the MoviPrep mixture, it is recommended that you drink 16 ounces of a clear liquid of your choice. This step helps to ensure your body remains hydrated during the bowel preparation process.

The day of your procedure

- Five hours before your colonoscopy begin to prepare the MoviPrep. Empty 1 pouch A and 1 pouch B into the disposable container. Add lukewarm drinking water to the top line of the container. Mix to dissolve.
- Start drinking the MoviPrep mixture. Drink an eight-ounce glass of the prep mixture every 15 minutes. You will drink a total of one liter or the entire container of the MoviPrep mixture. It will take you one hour to finish drinking it.
- After you finish the MoviPrep mixture, it is recommended that you drink 16 ounces of a clear liquid of your choice. This step helps to ensure your body remains hydrated during the bowel preparation process.
- You will need to be near a bathroom once you start drinking your prep mixture. You will have diarrhea and will need to be able to reach a bathroom quickly.
- Some people may feel nauseous, bloated, or vomit while drinking the bowel preparation. If this happens, take a 15-minute break, and then try drinking the prep mixture.

- Your stool should be no longer formed, but a clear or yellow liquid.
 - You may drink clear liquids until two hours before your check-in time.
 - Avoid red or purple liquids, dairy products, protein shakes, alcoholic beverages, chewing tobacco, and illicit drugs. The use of these will result in your procedure being canceled.
 - You may take your medications with a sip of water.
 - Remember, do not eat any solid foods.
 - You may want to allow extra time to travel to your appointment as you may need to stop and use a restroom along the way.
-

Notes / Other Instructions: